

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the spermatick Vessels; e, e, a Branch of the spermatick Vessels along the Side of the Intestin, compressed by the transversal Fibres, and inslected through its whole Extent in an uniform Manner; f, f, f, the Windings and Turnings of the spermatick Vessels; g, g, the transversal Fibres; h, the cartilaginous Membrane surrounding the Anus; i, the small Slit in its Middle; k, the very sine capillary Process; m, m, the small Bladders covering the Skin.

Fig. 7. A Male Worm inverted, and disected about the Anus, in Order to see with Ease the Duct lying under the alimentary Canal; a, the wrinkled alimentary Canal; b, the whitish Duct under the wrinkled Canal; c, the spermatick Vessels.

Fig. 8. a, the vermicular Process of the spermatick Vessels; b, b, the Branches of the spermatick Vessels, freed from their Windings; c, c, the same Branches dissected.

Bout five Years ago, I laid before the Society fome Observations made in an Ostrich dissected, by Order of Sir Hans Sleane, Bart. to which Account I beg Leave to add two or three more Observations which escaped my Notice in my former Dissection. And First, the Eye, the Figure of which, when Pp 2 taken

V. Some material Observations upon dissecting an Ostrich, made lately by Mr. Ranby, Surgeon to his Majesty's Houshold, and F. R. S.

taken out of the Orbit, I think particular, being almost triangular, with some litle Variation in the Bony Scales, from that in *Phil. Trans.* No. 386. The Contents of the Stomach were of such a Kind, that they were hardly capable (without very great Alteration) of

passing the lower Orifice, which is very small.

The Diameter of the *Duodenum* much smaller than any of the Intestines, and free from *Valves*, as are the *Jejunum* and *Ileum*, except the latter, which has a few *Valves*, as it approaches near the *Colon*. The *Colon* was uneven, with very regular Cells: These Cells were formed by *Valves*, which were on the Inside, and transversely situated, each making more than half a Circle.

The Parts in other Respects answer the Description given by the several curious Gentlementhat have diffected this Animal.